Reset Form | Print Form

FLORIDA MEDICAID PRIOR AUTHORIZATION

COLONY STIMULATING FACTORS

Preferred: Leukine®, Neupogen®, Nyvepria™

Clinical PA required (Non-Preferred): Fulphila™/Fylnetra®/Granix®/Neulasta®/ Nivestym®/Releuko®/Rolvedon™/Stimufend®/Udenyca®/Zarxio®/Ziextenzo™

Note: Form must be completed in full. An incomplete form may be returned.

Recipient's Medicaid ID #

Date of Birth (MM/DD/YYYY) / /

Recipient's Full Name

Prescriber's Full Name

Prescriber License # (ME, OS, ARNP, PA)

Prescriber Phone Number - -

Prescriber Fax Number - -

Pharmacy Name

Pharmacy Medicaid Provider #

Pharmacy Phone Number - -

Pharmacy Fax Number - -

Drug Name/Strength/NDC (if available) submitted on claim: _____

1. What is the diagnosis or the indication for the product? Please check below **AND** submit supporting documentation indicating the diagnosis.
   - ☐ Cancer patient receiving myelosuppressive chemotherapy
   - ☐ Cancer patient receiving bone marrow transplant
   - ☐ Patient receiving induction or consolidated chemotherapy for acute myeloid leukemia (AML)
   - ☐ Peripheral blood progenitor cell collection and therapy in cancer patient
   - ☐ Acute exposure to myelosuppressive doses of radiation in patient
   - ☐ Severe neutropenia in acquired immunodeficiency syndrome (AIDS) patient on antiretroviral therapy
   - ☐ Severe chronic neutropenia in patient (select from the following):
     - ☐ Congenital ☐ Cyclic ☐ Idiopathic

Mail or Fax Information to:
Magellan Medicaid Administration, Inc.
Prior Authorization
P. O. Box 7082
Tallahassee, FL 32314-7082
Phone: 877-553-7481
Fax: 877-614-1078

FLORIDA MEDICAID PRIOR AUTHORIZATION

COLONY STIMULATING FACTORS

Preferred: Leukine®, Neupogen®, Nyvepria™

Clinical PA required (Non-Preferred): Fulphila™/Fylnetra®/Granix®/Neulasta®/ Nivestym®/Releuko®/Rolvedon™/Stimufend®/Udenyca®/Zarxio®/Ziextenzo™

Note: Form must be completed in full. An incomplete form may be returned.

Recipient's Full Name

2. This is: ☐ New therapy **OR** ☐ Continuation of therapy
3. Can the prescriber attest the disease state or prescribed regimen is high risk (> 20%) for febrile neutropenia? ☐ Yes ☐ No
4. Lab test date: _____
   Absolute neutrophil count (ANC): _____ cells/mm^3
5. What is the date range of therapy? Begin date: _____ End date: _____
6. What will be the dosage and frequency of dosing? _____

Prescriber's Signature: _____ Date: _____

REQUIRED FOR REVIEW: Copies of medical records (i.e., diagnostic evaluations and recent chart notes) and the most recent copies of related labs. The provider must retain copies of all documentation for five years.

Mail or Fax Information to:
Magellan Medicaid Administration, Inc.
Prior Authorization
P. O. Box 7082
Tallahassee, FL 32314-7082
Phone: 877-553-7481
Fax: 877-614-1078

Approved Indications for Zarxio® and Nivestym®

- Cancer patients (Note that they do not have to meet ANC criteria. If they have the indication, approve.):
  - If patient has not yet undergone chemotherapy but it has been prescribed, no ANC is required
  - Cancer patients receiving myelosuppressive chemotherapy (approve up to 12 months)
  - Cancer patients receiving bone marrow transplants (approve up to 12 months)
  - Patient receiving induction or consolidated chemotherapy for AML (approve up to 12 months)
  - Peripheral blood progenitor cell collection and therapy in cancer patients (approve up to 12 months)
- Severe chronic neutropenia – ANC now required
  - All lab documentation must be on official lab letterhead – handwritten labs are not acceptable
  - The ANC is 1500 or less
  - Congenital, cyclic, or idiopathic (approve up to 12 months)
- AIDS – ANC required
  - Severe neutropenia in AIDS patients on antiretroviral therapy
  - Initial Therapy: ANC is 1000 or less
  - Continuation of Therapy: ANC is 1600 or less
  - All lab documentation must be on official lab letterhead – handwritten labs are not acceptable. (Approve for 6 months)

Approved Indications for Releuko®

- Cancer patients (Note that they do not have to meet ANC criteria. If they have the indication, approve.):
  - If patient has not yet undergone chemotherapy but it has been prescribed, no ANC is required
  - Cancer patients receiving myelosuppressive chemotherapy (approve up to 12 months)
  - Cancer patients receiving bone marrow transplants (approve up to 12 months)
  - Patient receiving induction or consolidated chemotherapy for AML (approve up to 12 months)
- Severe chronic neutropenia – ANC now required
  - All lab documentation must be on official lab letterhead – handwritten labs are not acceptable
  - The ANC is 1500 or less
  - Congenital, cyclic, or idiopathic (approve up to 12 months)

Approved Indications for Udenyca®, Neulasta®, Ziextenzo™, Fulphila™, Fylnetra®, Rolvedon™, and Stimufend®

- Chemotherapy-induced neutropenia
  - Cancer patient with non-myeloid malignancies receiving myelosuppressive chemotherapy (approve up to 12 months)
- Dosage: 6 mg subcutaneous once per chemotherapy cycle
- Patient acutely exposed to myelosuppressive doses of radiation (hematopoietic syndrome of acute radiation syndrome) (Neulasta® and Udenyca® only)
- Dosage: Two doses, 6 mg subcutaneous, each one week apart

Note:

- Do not administer in the period 14 days before and 24 hours after administration of cytotoxic chemotherapy.
- Documentation of the ANC and/or lab values is not required.
- Not indicated for severe chronic neutropenia.
- Not indicated for neutropenia associated with human immunodeficiency virus (HIV)/AIDS.

Approved Indications for Granix®

- Chemotherapy-induced neutropenia:
  - Cancer patient with non-myeloid malignancies receiving myelosuppressive chemotherapy (approve up to 12 months)
  - Dosage: 5 mcg/kg/day subcutaneously

Note:

- Do not administer in the period 24 hours before and 24 hours after administration of cytotoxic chemotherapy.
- Documentation of the ANC and/or lab values is not required.
- Not indicated for severe chronic neutropenia.
- Not indicated for neutropenia associated with HIV/AIDS.