



FLORIDA MEDICAID PRIOR AUTHORIZATION

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

Recipient's Medicaid ID#

Grid for Recipient's Medicaid ID#

Date of Birth (MM/DD/YYYY)

Grid for Date of Birth

Recipient's Full Name

Grid for Recipient's Full Name

Prescriber's Full Name

Grid for Prescriber's Full Name

Prescriber's NPI

Grid for Prescriber's NPI

Prescriber's Phone Number

Grid for Prescriber's Phone Number

Prescriber's Fax Number

Grid for Prescriber's Fax Number

PROVIDER TYPE OR SPECIALTY: CHILD UNDER STATE CARE/CUSTODY: PATIENT: MEDICATION REQUEST: HEIGHT: WEIGHT: BMI: *BMI %: BMI Calculator: *

1. Medication Requested:

Table with 4 columns: Requested Antipsychotic(s), Strength, Directions, Quantity

2. Diagnosis:

- ADHD, Disruptive Behavior Disorder, Disruptive Mood Dysregulation Disorder, Autism Spectrum, Schizophrenia, Other, Bipolar Disorder, Schizoaffective Disorder

3. Target Symptoms:

- Aggression, Impulsivity, Irritability, Self Injurious Behavior, Other

4. Severity of Target Symptoms:

- 1 Mild, 2 Moderate, 3 Marked, 4 Severe, 5 Extreme

5. Functional Impairment:

- 1 Mild, 2 Moderate, 3 Marked, 4 Severe, 5 Extreme

6. Previous Antipsychotic Trials in last 12 Months

Table with 4 columns: Antipsychotic Medication, Start Dates, End Dates, Maximum Dose (Per Day)



FLORIDA MEDICAID PRIOR AUTHORIZATION

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

Recipient's Full Name

Grid for recipient's full name

7. List all other psychotherapeutic medications the patient is taking concurrently with the antipsychotic (i.e., antidepressants, mood stabilizers, anxiolytics, etc.).

Table with 2 columns: Psychotherapeutic Medication, Dose/day

Table with 2 columns: Psychotherapeutic Medication, Dose/day

8. Rationale for prescribing antipsychotic above maximum recommended dose? (if applicable)

9. Is your intent to target lower dose antipsychotic treatment?

Yes No

10. Rationale for prescribing 2 or more antipsychotics for >60 days (if applicable):

11. If your request is for two antipsychotics:

Is the plan to cross taper, with antipsychotic monotherapy resumed within the next 60 days?

Yes No

If YES, please provide the cross taper plan:

12. Have metabolic monitoring labs* (fasting lipids and glucose) been performed within the last 6 months?

*Official lab results (most recent) must be attached. For continuation of therapy, labs are required.

Yes No Date:

13. Has an assessment for Tardive Dyskinesia (TD) been done in the last 6 months?

AIMs: Yes No

DISCUS: Yes No

Date:

*Official Form or notation (most recent) must be attached.

14. Monitoring Plan: RTC: Labs: q months TD Screen: q months

Labs: CBC Prolactin CMP Lipid Profile Other, specify:

15. Next Appointment Date:

Prescriber's Signature: Date:

REQUIRED FOR REVIEW: All copies of medical records (e.g., diagnostic evaluations and recent chart notes), and the most recent copies of related labs. The provider must retain copies of all documentation for five years.

Mail or Fax Information to: Magellan Medicaid Administration, Inc. Prior Authorization P. O. Box 7082 Tallahassee, FL 32314-7082 Phone: 877-553-7481 Fax: 877-614-1078



FLORIDA MEDICAID PRIOR AUTHORIZATION

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

FDA-approved agents and doses are considered most appropriate.

FDA Information for 6–17 Age Group		
Medication and Approved Use	Age Range	Dosing Instructions
Aripiprazole		
Bipolar Disorder (manic or mixed episodes)	Pediatric age 10–17	Initial dose: 2 mg/day Recommended dose: 10 mg/day Maximum dose: 30 mg/day
Schizophrenia	Pediatric age 13–17	Initial dose: 2 mg/day Recommended dose: 10 mg/day Maximum dose: 30 mg/day
Irritability associated with Autism	Pediatric age 6–17	Initial dose: 2 mg/day Recommended dose: 5–10 mg/day Maximum dose: 15 mg/day
Lurasidone		
Bipolar I Disorder (depression)	Pediatric age 10-17	Initial dose: 20 mg/day Recommended dose: 20-80 mg/day Maximum dose: 80 mg/day
Schizophrenia	Pediatric age 10-17	Initial dose: 40 mg/day Recommended dose: 40-80 mg/day Maximum dose: 80 mg/day
Olanzapine		
Bipolar I Disorder (manic or mixed episodes)	Pediatric age 13–17	Oral Formulation Initial dose: 2.5–5 mg/day Target dose: 10 mg/day
Schizophrenia	Pediatric age 13–17	Initial dose: 2.5–5 mg/day Target dose: 10 mg/day
Paliperidone		
Schizophrenia	Pediatric age 12–17	Weight < 51kg: Initial Dose (3 mg/day) Recommended Dose (3–6 mg/day) Maximum Dose (6 mg/day) Weight ≥ 51kg: Initial Dose (3 mg/day) Recommended Dose (3–12 mg/day) Maximum Dose (12 mg/day)
Risperidone		
Bipolar I Disorder (manic or mixed episodes)	Pediatric age 10–17	Initial dose: 0.5 mg/day Titration: 0.5–1 mg/day Recommended dose: 2.5 mg/day Effective dose range: 0.5–6 mg/day

Confidentiality Notice: The documents accompanying this transmission contain confidential health information that is legally privileged. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender (via return fax) immediately and arrange for the return or destruction of these documents. Distribution, reproduction or any other use of this transmission by any party other than the intended recipient is strictly prohibited.



FLORIDA MEDICAID PRIOR AUTHORIZATION

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

FDA Information for 6–17 Age Group		
Medication and Approved Use	Age Range	Dosing Instructions
Irritability associated with Autism	Pediatric age 5–16	<p>Initial dose: 0.25 mg/day (< 20 kg); 0.5 mg/day (≥ 20 kg)</p> <p>Titration: 0.25–0.5 mg at > or = 2 weeks</p> <p>Recommended dose: 0.5 mg/day (< 20 kg); 1 mg/day (≥ 20 kg)</p> <p>Effective dose range: 0.5–3 mg/day</p>
Schizophrenia	Pediatric age 13–17	<p>Initial dose: 0.5 mg/day</p> <p>Titration: 0.5–1 mg/day</p> <p>Target dose: 3 mg/day</p> <p>Effective dose range: 1–6 mg/day</p>
Quetiapine		
Bipolar I Disorder (mania)	Pediatric age 10–17	<p>Information provided is for the immediate release table formulation</p> <p>Day 1: 25 mg twice a day</p> <p>Day 2: Twice daily dosing totaling 100 mg</p> <p>Day 3: Twice daily dosing totaling 200 mg</p> <p>Day 4: Twice daily dosing totaling 300 mg</p> <p>Day 5: Twice daily dosing totaling 400 mg</p> <p>Further adjustments should be in increments no greater than 100 mg/per day within the recommended dose range of 400–600 mg/per day. Based on response and tolerability, may be administered three times daily.</p>
Schizophrenia	Pediatric age 12–17	<p>Information provided is for the immediate release tablet formulation</p> <p>Day 1: 25 mg twice daily</p> <p>Day 2: Twice daily dosing totaling 100 mg</p> <p>Day 3: Twice daily dosing totaling 200 mg</p> <p>Day 4: Twice daily dosing totaling 300 mg</p> <p>Day 5: Twice daily dosing totaling 400 mg</p> <p>Recommend dose range: 400–800 mg/day</p> <p>Further adjustments should be in increments no greater than 100 mg/per day within the recommended dose range of 400–800 mg/per day. Based on response and tolerability, may be administered three times daily.</p>

Helpful Links:

- Access the following information at <http://floridabhcenter.org/index.html>:
 - **Antipsychotic High Dosing Table for Children and Adolescents**
 - **AIMS/DISCUS forms**
 - Florida Medicaid **Best Practice Psychotherapeutic Medication Guidelines for Children and Adolescents**
- The Centers for Disease Control and Prevention (CDC) **BMI Calculator for Children and Teens:** <https://www.cdc.gov/healthyweight/bmi/calculator.html>

Confidentiality Notice: The documents accompanying this transmission contain confidential health information that is legally privileged. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender (via return fax) immediately and arrange for the return or destruction of these documents. Distribution, reproduction or any other use of this transmission by any party other than the intended recipient is strictly prohibited.