

Florida Medicaid Cesarean Section Rate Calculation Specifications 2019

Description provider) that were delivered via cesarean section (C-section). January 1, 2019 January 1, 2019 Initial Measurement Period Plans must use this measurement period to calculate the C-section rate to determine which Identified Providers are qualified to receive the incentive payment as of October 1, 2020. July 1, 2020 – November 30, 2020 Plans may use this re-measurement period to calculate the C-section rate to determine which Identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a delivery via a cesarean section (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the follo	_	The percentage of single liveborn Medicaid births in a practice (pay to		
Initial Measurement Period January 1, 2019 – December 31, 2019 Initial Measurement Period Plans must use this measurement period to calculate the C-section rate to determine which Identified Providers are qualified to receive the incentive payment as of October 1, 2020. July 1, 2020 – November 30, 2020 Re-measurement Period Plans may use this re-measurement period to calculate the C-section rate to determine which Identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Numerator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Denominator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient bad a single liveborn delivery (use codes in Table 1). • Re	Description			
Period Plans must use this measurement period to calculate the C-section rate to determine which identified Providers are qualified to receive the incentive payment as of October 1, 2020. July 1, 2020 – November 30, 2020 Plans may use this re-measurement period to calculate the C-section rate to determine which identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider revel. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Plans must exclude births that have a diagnosis code listed in Table 1). Recipient had a single liveborn delivery (use codes in Table 1). Plans must exclude births that have a diagnosis code listed in the measure denominator: Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between t				
Period Plans must use this measurement period to calculate the C-section rate to determine which identified Providers are qualified to receive the incentive payment as of October 1, 2020. July 1, 2020 – November 30, 2020 Plans may use this re-measurement period to calculate the C-section rate to determine which identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider revel. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Plans must exclude births that have a diagnosis code listed in Table 1). Recipient had a single liveborn delivery (use codes in Table 1). Plans must exclude births that have a diagnosis code listed in the measure denominator: Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between t	Initial Measurement			
Determine which identified Providers are qualified to receive the incentive payment as of October 1, 2020. Re-measurement Period July 1, 2020 – November 30, 2020 Plans may use this re-measurement period to calculate the C-section rate to determine which Identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Denominator Recipient's baby was delivered by a Provider level. Denominator Recipient's baby was delivered by a Provider level. Denominator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient ad a delivery via a vaginal or cesarean section (use codes in Table 1				
Re-measurement Period July 1, 2020 – November 30, 2020 Plans may use this re-measurement period to calculate the C-section rate to determine which Identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a single liveborn delivery (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Denominator Recipient had a delivery via a vaginal or cesarean section (use codes in Table 4). Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plenominator Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider who had a delivery date of service during the measurement period (i chou	•		
Re-measurement Period Plans may use this re-measurement period to calculate the C-section rate to determine which Identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Plans must exclude births that have a diagnosis code listed in Table 1). Recipient had a single liveborn delivery (use codes in Table 1). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Denominator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator Recipient ad a single liveborn delivery (use codes in Table 1). Recipient ad a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level.				
Recineasurement Period determine which Identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Numerator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Denominator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Plans must exclude births that have a diagnosis code listed in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). Becipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that h		July 1, 2020 – November 30, 2020		
Recineasurement Period determine which Identified Providers are newly qualified to receive the incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Numerator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Denominator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Plans must exclude births that have a diagnosis code listed in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). Becipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that h		Plans may use this re-measurement period to calculate the C-section rate to		
Period incentive payment as of April 1, 2021. The re-measurement period cannot be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Numerator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The numer of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Denominator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4				
Denominator be used to remove an October 1, 2020 qualified provider from receiving the April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2	Period			
April 1, 2021 incentive payment. The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2). • Recipient had a de				
Denominator The number of unduplicated Medicaid recipients between the ages of 10 and 60 who meet each of the following criteria is included in the measure numerator: Numerator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The denominator should be calculated at the practice (pay to provider) level, rather than at the render				
60 who meet each of the following criteria is included in the measure numerator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level.				
Numerator: Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient ad a single liveborn delivery (use codes in Table 1). Recipient ad a single liveborn delivery (use codes in Table 1). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level.				
Numerator of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Calculation Numerator * 100		•		
Numerator of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Calculation Numerator * 100		Becipient's haby was delivered by a Provider who had a delivery date		
Numerator for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator				
Numerator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Table 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Calculation Numerator				
 Recipient had a delivery via a cesarean section (use codes in Table 2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. 	Numerator			
2). Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Calculation Numerator * 100	Numerator			
Plans must exclude births that have a diagnosis code listed in Table 4. The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100				
Denominator The numerator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100				
rather than at the rendering/treating provider level. The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator		Plans must exclude births that have a diagnosis code listed in Table 4.		
Denominator The number of unduplicated Medicaid recipients between the ages of 10-60 who meet each of the following criteria is included in the measure denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator		The numerator should be calculated at the practice (pay to provider) level,		
who meet each of the following criteria is included in the measure denominator: Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100		rather than at the rendering/treating provider level.		
denominator: • Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Denominator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100				
 Recipient's baby was delivered by a Provider who had a delivery date of service during the measurement period (see above for date spans for each period). Recipient had a single liveborn delivery (use codes in Table 1). Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100 				
Denominator of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100		denominator:		
Denominator of service during the measurement period (see above for date spans for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100		Recipient's baby was delivered by a Provider who had a delivery date		
Denominator for each period). • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100				
Denominator • Recipient had a single liveborn delivery (use codes in Table 1). • Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100				
Recipient had a delivery via a vaginal or cesarean section (use codes in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. <u>Numerator</u> * 100	Denominator			
in Tables 2 and 3). Plans must exclude births that have a diagnosis code listed in Table 4. The denominator should be calculated at the practice (pay to provider) level, rather than at the rendering/treating provider level. Numerator * 100				
Calculation Numerator Numerator * 100				
calculation Numerator * 100		Plans must exclude births that have a diagnosis code listed in Table 4.		
calculation Numerator * 100		The denominator should be calculated at the practice (pay to provider) level,		
Calculation Numerator * 100		rather than at the rendering/treating provider level.		
	Colouistion	Numerator + 100		
	Calculation	Denominator ^ 100		



Codes used to Identify Included Births

 Table 1: ICD-9 and ICD-10 Diagnosis Codes for identifying Singleton Liveborn

ICD-9	ICD-10	Description
V27.0	Z370	Single live-born

Table 2: CPT Procedure Codes for Identifying Cesarean Section Deliveries

CPT Procedure Codes	CPT Procedure Code Description
59510	Global code: routine obstetric care including
	antepartum care, C-section delivery, and
	postpartum
59514	C-section delivery only
59515	C-section delivery including postpartum care
59618	Routine obstetric care including antepartum care, cesarean delivery, and postpartum care. Following an attempted vaginal delivery after previous C- section delivery.
59525	C-section delivery with removal of uterus (hysterectomy)
59620	Cesarean delivery only, following attempted vaginal delivery after previous cesarean delivery
59622	C-section delivery (following attempted vaginal delivery after previous C-section delivery; including postpartum care
540	APR – DRG Inpatient C-Section delivery, liveborn

 Table 3: CPT Procedure Codes for Identifying Vaginal Deliveries

CPT Procedure Codes	CPT Procedure Code Description
59400	Routine obstetric care including antepartum care,
	vaginal delivery (with or without episiotomy, and/or
	forceps) and postpartum care
59409	Vaginal delivery only (with or without episiotomy
	and/or forceps)
59410	Vaginal delivery only (with or without episiotomy
	and/or forceps); including postpartum care
59614	Vaginal delivery only, after previous cesarean
	delivery (with or without episiotomy and/or forceps);
	including postpartum care
59610	Routine obstetric care including antepartum care,
	vaginal delivery (with or without episiotomy, and/or
	forceps) and postpartum care, after previous
	cesarean delivery
APR-DRG Codes	APR-DRG Desc
541	VAGINAL DELIVERY W STERILIZATION &/OR
	D&C
542	VAGINAL DELIVERY W COMPLICATING
	PROCEDURES EXC STERILIZATION &/OR D&C
560	VAGINAL DELIVERY



Codes Used to Identify Excluded Births Table 4: ICD-9 and ICD-10 Diagnosis Codes for identifying Stillborn and Multiple Gestation Births

ICD- 9	ICD-10	Description
V27.1	Z37.1	Outcome of delivery, single stillborn
V27.2	Z37.2	Multiple gestations
V27.3	Z37.3	Twins, one live-born and one stillborn
V27.4	Z37.4	Twins, both stillborn
V27.5	Z37.59	Other multiple births, all live-born
V27.6	Z37.69	Other multiple births, some live-born
V27.7	Z37.7	Twins, both live-born
656.40-656.43	O36.4XXØ	Stillborn or intrauterine death
	O3Ø.ØØ9	Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, unspecified trimester
651.00 - 651.93	O3Ø.91	Multiple gestations, unspecified first, second, or third trimester
	O3Ø.92 O3Ø.93	